

529245

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
8 April 2004 (08.04.2004)

PCT

(10) International Publication Number  
**WO 2004/028281 A1**

(51) International Patent Classification<sup>7</sup>: A23P 1/16, A23L 1/05, A23J 3/00, A23G 3/00, 1/00, A21D 13/00, A23L 1/053, 1/054, 1/056, 1/0534, A61K 7/00

(21) International Application Number:  
PCT/EP2003/010740

(22) International Filing Date:  
26 September 2003 (26.09.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
02021653.7 27 September 2002 (27.09.2002) EP

(71) Applicant (for all designated States except US): NESTEC S.A., A SWISS BODY CORPORATE OF AVENUE NESTLE 55 [CH/CH]; CH-1800 Vevey (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): KOLODZIEJCZYK, Eric [FR/CH]; Rue de la Confrérie 8, CH-1800 Vevey (CH). SCHMITT, Christophe [FR/CH]; Route d'Arnier 16, CH-1092 Belmont S/Lausanne (CH).

(74) Agent: THOMAS, Alain; Avenue Mestlé 55, CH-1800 Vevey (CH).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES,

[Continued on next page]

(54) Title: INTERFACE STABILISATION OF A PRODUCT WITH 2 OR MORE PHASES WITH A PROTEIN-POLYSACCHARIDE COMPLEX



(57) Abstract: The present invention concerns a product taken from the group consisting of a foam, an emulsion, a foamed emulsion, a dispersed emulsion and a foamed dispersion, wherein the interface water-air, water-oil or water-solid comprises a complex formed instantaneously by the mixture of at least a protein (or peptide) and at least a polysaccharide oppositely charged or the mixture of two proteins oppositely charged, said product being in a pH range within which the electrostatic interaction between both compounds oppositely charged occurs and wherein the total amount of protein and polysaccharide is comprised between 0.01 and 5 % in weight.

WO 2004/028281 A1



FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY,

TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

- of inventorship (Rule 4.17(iv)) for US only

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERFACE STABILISATION OF A PRODUCT WITH 2 OR MORE PHASES WITH A  
PROTEIN-POLYSACCHARIDE COMPLEX

The present invention concerns a two-phases or more product with enhanced stability.

5

A two-phases product, like a foamed product are very known on the market and appreciated by the consumer. An emulsion, like a mayonnaise, is also very spread on the market. For stabilizing the emulsion, an emulsifier is normally used, which is directly present in the bulk phase. The main drawback of this solution is the limitation of the diffusion of the emulsifier from the bulk to the interface resulting in a decrease in the final product stability. In the case of a foam, the control and design of texture is mainly achieved by adjusting the viscosity properties of the liquid bulk phase surrounding air bubbles (Walstra P. and De Roos A.L. (1993), Food Rev. Int., 9,503-525). To overcome the foam formation and foam stability problems, one generally combine surfactant molecules (phospholipids, fatty acids) together with tension active molecules (proteins). The former will initially decrease the interfacial area by adsorption at the interface, resulting in a high foam capacity. By contrast, the latter will form a viscoelastic layer around the bubbles, decreasing then the surface tension. This results in a higher stability of the foam. However, this combination has some drawbacks, since it requires the use of complex mixtures of surfactant and tension-active molecules. Moreover, it has been shown that both types of molecules are generally incompatible at the interface leading to interfacial phase separation and destabilisation (Mackie A.R. and al. (1999), J. Colloid Interf. Sc., 210, 157-166).

10

15

20

25

30

CONFIRMATION COPY

It is also possible to use protein-polysaccharide complexes to stabilize interfaces. This is the case for the US Patent No. 6'197'319, wherein a protein-polysaccharide complex is incorporated in a cosmetic composition, which is an emulsion. In this case, the complex is previously formed and then incorporated in the bulk. This is the same for the EP Patent No. 340'035 : a microfragmented ionic polysaccharide/protein complex dispersion is formed to be used as fat substitute in food products, such as ice cream, salad dressings, dips, spreads and sauces. The US Patent No. 3'944'680 concerns a process for the preparation of an aqueous oil emulsion of prolonged storage life. In this case, the complex protein/polysaccharide is formed in the bulk, and there is a problem of diffusion of the complex from the bulk to the interface as already mentioned in the preceding paragraph.

Hence, these protein-polysaccharide complexes have been shown to form upon electrostatic attraction in well defined conditions of pH, ionic strength, protein to polysaccharide ratio, total biopolymer concentration, temperature or pressure (Schmitt C. *et al.* (1998), *Crit. Rev. Food Sci. Nutr.*, 38, 689-753). In addition, various studies have demonstrated that these complexes exhibited better functional properties such as gelation, emulsification and foaming than that of the biopolymers alone. However, it is also known that the formation of complexes through electrostatic attraction between protein and polysaccharide lead to the associative phase separation phenomena (Piculell L. and Lindman B. (1992), *Adv. Colloid Interf. Sci.*, 41, 149-178; Doublier J.-L. *et*

al. (2000), *Curr. Opinion Colloid Interf. Sci.*, 5, 202-214) or complex coacervation (Bungenberg de Jong H.G. (1936), *La coacervation complexe et son importance en biologie*, E. Fauré-Fremiet Ed, vol 1, Paris: Hermann et Cie). During associative phase separation which is a time dependent mechanism, the initial electrostatic protein-polysaccharide complexes interact ones with the others because of charge neutralisation to increase the electrostatic entropy of the system by release of counter-ions in the medium (Tolstoguzov V.B. (1997), *Protein-polysaccharide interactions*, S. Damodaran and A. Paraf Eds, *Food Proteins and their Applications*, pp 171-198, New York: Marcel Dekker Inc). Coming closer to the thermodynamic equilibrium, the complexes become insoluble and form liquid droplets called coacervates. These coacervates finally form a concentrated liquid phase at equilibrium with a very diluted phase containing mainly solvent (Mattisson K.W. et al. (1999), *Macromol. Symp.*, 140, 53-76). The size of the successive entities formed ranged from tens of nanometers for the initial macromolecular complexes (Xia J. (1993) *Macromolecules*, 26, 6688-6690; Bowman W. (1997), *Macromolecules*, 30, 3262-3270) to hundreds of microns for the coacervates (Schmitt C. et al. (2001), *Colloids and Surf. B: Biointerf.*, 20, 267-280). In terms of interfacial activity, it is well know that the coefficient of diffusion of the surface-active components is very important. Large molecular weight surface-active components (for example, protein-polysaccharide complexes) go very slowly at the interface to the contrary of low molecular weight surface-active components (sugar-esters, triglycerides). However, the former are much more effective for interfacial



stabilisation (Dickinson E. and Galazka V.B. (1991), Food Hydrocolloids, 5, 281-296).

5 It has been demonstrated that protein/anionic polysaccharide or basic protein/acidic protein mixtures are able to improve significantly the foaming properties as compared to the protein taken alone (Ahmed & Dickinson, 1991, Food Hydrocolloids, pages 395-402; Poole, 1989, International Journal of Food Science and 10 Technology, pages 121-137 and GB 2134117 and GB 2179043). In the latter documents cited, no improvement of the foam stability is clearly described after mixing of the protein with another protein or polysaccharide. Thus, these papers refer only to increased foamability, which 15 did not foresee any improvement of stability. An important point that is not mentioned in these documents is the effect of the time on the surface properties of the formed complexes. Since complexation is mainly due to electrostatic interactions between the two compounds, a 20 charge neutralisation of the complexes arises with time (whatever the initial mixing ratio). It follows that more and more complexes are able to interact one with the other (no repulsion between the complexes), leading to an increase of the size of the complexes and to their 25 progressive insolubilisation. These two phenomena are very detrimental for the stabilisation of an interface since they reduce the capability of the complexes to remain in solution and to migrate at the interface. This is the main reason why the industrial use of 30 electrostatic complexes is scarce. Our invention allows circumventing these critical points in the use of protein/polysaccharide or protein/protein complexes. Since we are creating the complexes concomitantly with

the interface (gas/liquid, gas/solid, or solid/liquid), they remain soluble and are small enough to go at the interface (with the help of the energy input within the system). Once at the interface, these complexes rearrange  
5 to form coacervates that effectively form a stabilising film and that can be detected at the interface within finished food products by means of microscopic histochemical techniques.

10

According to the present invention, it is possible to stabilise efficiently interfaces using protein-polysaccharide or protein-protein electrostatic complexes formed at the same time than the interface they have to  
15 stabilise, because they have smaller size and a higher diffusion coefficient. In this case, the protein-polysaccharide complexes will be soluble and with sufficiently low molecular weight to be at the interface. Based on these observations, the objective of  
20 the present invention is, in the case of a at least two phases product, to control the surface properties between said phases.

25

The present invention concerns a product taken from the group consisting of a foam, an emulsion, a foamed emulsion, a dispersed emulsion and a foamed dispersion, wherein the interface water-air, water-oil or water-solid comprises a complex formed instantaneously at said  
30 interface by the mixture of at least a protein (or peptide) and at least a polysaccharide oppositely charged or the mixture of two proteins oppositely charged, said product being in a pH range within which the electrostatic interaction between both compounds

oppositely charged occurs and wherein the total amount of protein and polysaccharide is comprised between 0.01 and 5 % in weight.

5 The complex (or coacervate) in the product of the invention is formed instantaneously, directly during the preparation of said product and positioned directly at the interface water-air, water-oil and water-solid. The active compounds are in aqueous solutions or oil-in-water  
10 emulsions.

The purpose of the invention is to control the surface properties of the interfaces by using an ingredient mix of protein and polysaccharide or a mix of two proteins.  
15 The mix significantly enhances, in the case of foaming, the foam capacity, that means more foam is obtained and the foam stability (smaller air bubbles, less drainage). The mix can also be used for emulsions, and other at least two phases products.

20 The mix efficiency remains in the formation of electrostatic complexes under well defined conditions of pH (when electrostatic interaction occurs), temperature (from 0°C to room temperature), with a ratio protein to  
25 polysaccharide or ratio of both proteins from 1:20 to 20:1 and total biopolymer concentration between 0.01 and 5 % in weight. Under biopolymer, we understand the addition of the weight concentrations of protein and polysaccharide.

30 By using the complex in liquid form, in case of foam, it is possible to enhance the foam formation since the initially formed complex acts as surfactant. After the



foam is formed, the complex further interacts one with the other to form the so-called coacervates. These coacervates exhibit high viscoelastic properties and by the way rearrange at the interface to form a viscoelastic film that stabilise the foam.

A product containing protein-polysaccharide or protein-protein complex obtained through ionic interaction is used to produce different types of interfaces from liquid dispersion. In a first step, the molecular complex acts as surfactant, so that the interfacial area is decreased. In a second step, the complex rearrange at the interface in order to form coacervates that spread around the air bubbles (or oil droplets) , forming a viscoelastic film that stabilises the bubbles (or oils droplets) against destabilisation.

The instant formation of the protein-polysaccharide or protein-protein complex results in a characteristic structural signature combining both a typical structure and distribution of fats and a typical structure and composition of viscoelastic films at the water/air or water/oil or water/solid interfaces. This can be demonstrated with various microscopy techniques, in particular, by showing specifically the location of the protein(s) and the polysaccharide components within the interfaces: see below figures 5 and 6.

In the case of a foamed product, the active compounds are in aqueous form, and the interface concerned is an air-water interface. In the case of an emulsion, the active compounds are in an aqueous phase, and the interface concerned is an oil-water interface. In the case of a

foamed emulsion, both air-water and oil-water interfaces are concerned. In the case of a dispersed emulsion, both air-solid and oil-water interfaces are concerned. In the case of a foamed dispersion, both air-solid and water-air  
5 interfaces are concerned.

The protein of the product is taken from the group consisting of a protein from milk, soy, egg, meat, fish and plant. Under plant, we understand mainly cereals but  
10 also leguminosea . Most preferably, the protein is  $\beta$ -lactoglobulin, gelatin,  $\alpha$ -lactalbumin, bovin serum albumin, soy globulin, soy protein, wheat protein, whey protein, soy protein, egg white protein.

15 The polysaccharide is taken from the group consisting of charged natural or synthetic polysaccharides. Most preferably, the polysaccharide is acacia gum, carboxymethyl-cellulose, chitosan, xanthan, alginate, propylene-glycol alginate, carrageenans, low or high methoxylated  
20 pectins, arabinogalactans, rye arabinoxylans (Ax rye), wheat arabinoxylans (Ax wheat).

The product of the invention can be used either per se or in mixture with another product. In the case of a use in  
25 the food area, the final product is ice cream, a culinary product, chocolate, dessert, a dairy product , wafers, sponge cakes or a petfood product. It is also possible to use it in coffee creamer or dairy coffee creamer. In this case, the product of the invention is present in an  
30 amount of 10 to 100 % of the final product in weight.

In the case of a use in the cosmetic or perfume area, the product is used in an amount comprised between 10 and 100 % of the final product in weight.

5 The invention concerns further the processes for the preparation of the product of the invention. There are different possible products and the following description will consider all these possible ways.

10 In the case of a preparation of a foam product, the way of preparation is following :

a solution or a bulk mass of the at least one protein and a solution or a bulk mass of the at least one polysaccharide or a solution or a bulk mass of one  
15 protein and another solution or a bulk mass of one protein is injected with the air in a bulk mass or directly in the air in the case of two bulk masses using the multi-tube reactor described in Fig 1. The bulk mass can be a dairy product, containing sugar or not,  
20 containing a living organism or not. In this case, the polysaccharide is preferably acacia gum and the protein is  $\beta$ -lactoglobulin. The product has a final pH of around 4.2. The concentration of the protein and polysaccharide is around 0.01 to 5 % in weight. The ratio of protein to  
25 polysaccharide is around 20:1 to 1:20. The preparation is carried out at a temperature of around 4 to 50°C. In this case, the product obtained is the final product. The quantity of air injected is not critical and can vary between 10 and 700 % of the product.

30

According to a second embodiment of the preparation of a foam product, the way of preparation is following : a solution of at least one protein and a solution of at

least one polysaccharide or a solution of one protein and another solution of one protein are mixed together in the presence of air. This is the basic concept of the invention : the mixing of both active compounds creates  
5 per se the formation of the foam. If necessary, it is also possible to carry out a subsequent whipping. According to this way of proceeding, the product obtained is not the final product. It is then mixed with a preparation for ice cream, or with a preparation for a  
10 wafer.

In the case of a preparation of an emulsion the way of preparation is following :

A first part of an emulsion is stabilised with at least  
15 one protein, a second part of a second emulsion is stabilised with at least one polysaccharide or a second protein, and both emulsions are mixed together.

The lipid used for the first and the second  
emulsion is preferably palm oil, palm kernel, sunflower,  
20 safflower or olive oil or butterfat or butterfat or any of their mixtures. The main interest of this embodiment is the preparation of a mayonnaise, for example an egg-yolk free or low-fat mayonnaise. In this case, the protein is preferably  $\beta$ -lactoglobulin and the polysaccharide is  
25 acacia gum. The product has a final pH of around 4.2. The concentration of the protein and polysaccharide is around 0.01 to 5 % in weight. The preparation is carried out at a temperature at which the fat is liquid. In this case, the product obtained is the final product. This is a non  
30 foamed product.

It is also possible, according to a second embodiment of the preparation of an emulsion to have a foamed emulsion.

In this case, a solution of at least one protein and a solution of at least one polysaccharide or a solution of one protein and another solution of one protein are mixed together in the presence of air. This foamed product is  
5 then incorporated in an emulsion to obtain for example a foamed mayonnaise. In this case , the protein is  $\beta$ -lactoglobulin, the polysaccharide is acacia gum and the lipid phase is sunflower oil or olive oil. The overrun of whipping is around 10 to 700 %.

10

According to a further embodiment of the invention, it is possible to prepare directly a foamed emulsion. In this case, the way to proceed is following :

a bulk product is prepared using a first part of an  
15 emulsion which is stabilised with at least one protein, a second part of an emulsion is stabilised with at least one polysaccharide or a second protein, both emulsions are mixed together and diluted in the bulk product, then  
20 a new dispersion of the protein with a new dispersion of the polysaccharide are injected with air (using the foaming device described on Fig 1) in the bulk product to form the foamed bulk product.

In this case, the bulk product is preferably a mix for  
25 ice cream. The lipid used for the emulsion is preferably sunflower oil, palm oil, palm kernel oil or milk fat. The protein used is preferably  $\beta$ -lactoglobulin and the polysaccharide is acacia gum. The amount of both protein and polysaccharide is around 0.01 to 5 % in weight. The  
30 pH is around 4.2. The final product is then either frozen by a static freezing or by a dynamic freezing.



According to a further embodiment, a dispersed emulsion can be prepared. The way of preparation is following :  
a first part of an emulsion is stabilised with at least one protein, a second part of an emulsion is stabilised  
5 with the at least one polysaccharide or a second protein, and both emulsions are mixed together, the obtained final emulsion being then mixed with a base comprising particles. In a preferred embodiment, the base comprising particles is a chocolate base containing for  
10 example sugar, cocoa particles, milk powder, lecithine and other aroma.

The way of producing is following :  
A first part of an emulsion is stabilised with at least  
15 one protein, a second part of a second emulsion is stabilised with at least one polysaccharide or a second protein, and both emulsions are mixed together.  
The lipid used for the first and the second  
emulsion is preferably palm oil, palm kernel, sunflower,  
20 safflower or olive oil or butterfat or butterfat or any of their mixtures. The main interest of this embodiment is the preparation of a mayonnaise, for example an egg-yolk free or low-fat mayonnaise. In this case, the protein is preferably  $\beta$ -lactoglobulin (BLG) and the  
25 polysaccharide is acacia gum (AG). The product has a final pH of around 4.2. The concentration of the protein and polysaccharide is around 0.01 to 5 % in weight. The preparation is carried out at a temperature at which the fat is liquid. This emulsion product is then mixed with a  
30 bulk composed by lecithin, sugar and cocoa particles to obtain a chocolate base.

According to a last embodiment of the invention, a foamed dispersion is prepared. The way of producing is following :

5 a solution of the at least one protein and a solution of the at least one polysaccharide or a solution of one protein and another solution of one protein is injected with the air in a bulk of dispersed particles.

10 A typical example of a foamed dispersion is the preparation of a sorbet. In this case, a solution of the at least one protein and a solution of the at least one polysaccharide or a solution of one protein and another solution of one protein is injected with the air in a bulk mass. In this case , the bulk mass does not contain  
15 fat, but more a fruit, or juice or fruit puree.

The bulk dispersed particles can be a yoghurt preparation, or another acidic gel with bacteria or with a gel obtained by chemical acidification or heat  
20 treatment of gelling polysaccharides or proteins.

It is also possible to proceed according to a second way : a solution of at least one protein and a solution of at least one polysaccharide or a solution of one protein and  
25 another solution of one protein are mixed together in the presence of air. The bulk of dispersed particles is then mixed with this foam to obtain the foamed dispersion.

For this last embodiment, it is possible to consider the  
30 preparation of wafers or caramel.

The invention concerns further the device for carrying out the process of the invention. This device comprises on a frame :

- a first pipe for the arrival of an emulsion or dispersion with a protein
- a second pipe for the arrival of the gas
- a third pipe for the arrival of an emulsion or dispersion with a polysaccharide,

these three pipes arriving on a main channel, being disposed perpendicular and staggered along said main channel , the first pipe forming the central pipe on the main channel, the second pipe forming the intermediate pipe on the main channel and the third pipe forming the external pipe on the main channel and wherein the outlet on the main channel of the central and the intermediate pipe are staggered.

A further object of the present invention is to have a product, wherein the stability of its structure during storage or heat shock is kept practically unchanged. Under storage in the present description, we understand a storage of one to twelve months. Considering the case of ice cream, by practically unchanged, we understand that the volume of air bubbles does not decrease much by taking out and putting again said ice cream several time in the freezer. The consumer has therefore a product, wherein the quality remains practically unchanged during its whole storage. This is the same for the other products considered in the present patent .

30

A further advantage of the present invention is that in the case of a co-emulsion, no emulsifier is needed in the product according to the invention.

The description is now made in relation with the figures, wherein

- 5 Figure 1 shows a perspective view of the device (triple needle) according to the invention and

Figure 2 is an enlarged view of the outlet of the device.

- 10 Figure 3 is a microscopical view of the air bubbles in the ice-cream according to example 2 further to their dispersion in the medium used for the quantitative analysis,

- 15 Figure 4 is a graph of the cumulative volume distribution of air bubbles analysed in example 2,

Figure 5 show confocal scanning laser microscopy of products of example 2,

20

Figure 6 show confocal scanning laser microscopy of products of example 2, at a higher resolution and

- 25 Figure 7 shows the interface of an air bubble where coacervates were formed instantaneously at the interface in a model system of an air bubble in a biopolymer mixture solution.

- 30 The injection system is on a frame (4). In this frame, there are 3 pipes. The pipe (1) is used for the arrival of the first dispersion with the protein, the pipe (2) is used for the entrance of the gas, that is the air and the pipe (3) is used for the arrival of the second dispersion

with the polysaccharide. The 3 pipes of the device arrive on a main channel (8) : they are disposed perpendicularly and staggered along said main channel (8). This device is used for the different examples in the specification and hereunder.

The important feature is that at the point of the mixing of the gas with the first and the second dispersion the outlet of pipes (5) and (6) are staggered as shown on figure 2. This allows a reaction time between the components of the mixture.

Figure 3 is the air bubbles dispersion found after whipping to 100% overrun at +4°C the ice-cream recipe described in example 2 using a Mondomix apparatus. Picture A is a classical mixture (10% non-fat milk solids) that did not contain the  $\beta$ -lactoglobulin and the acacia gum. The shearing speed was set to 800 rpm. Picture B is the same recipe but part of the non-fat milk solids was replaced by 2.5% whey protein isolate and 1.25% acacia gum. Here also shearing speed was 800 rpm. Picture C was taken after aeration of the recipes described in example 2 using the injection device (Fig. 1) coupled with the whipping device (Mondomix). The air bubble size is the same than in A and B, but the shearing rate is only 400 rpm. These results clearly demonstrated the highest foamability of the ice cream recipe containing protein-polysaccharide complexes instantaneously produced at the interface.

The air bubble analysis was carried out according to the procedure briefly described below :



An aliquot of the product is weighed and dispersed in a medium of high viscosity. The composition of the dispersing medium is designed to stabilize the air bubbles. The dispersion is submitted to an automated  
5 quantitative image analysis. The same procedure is taken for figure 4.

Figure 4 represents the cumulated volume size distributions of the recipes corresponding to the  
10 previous control (D), BLG/AG control (E) and triple needle BLG/AG (F) recipes described before (A) and after (B) heat shock treatment. The graphs represent the cumulative distribution of air bubbles in percent of the total volume of air analysed (x axis gives the air bubble  
15 diameter in mm and y axis gives in % the cumulated air volume analysed). Samples were equilibrated for one week at  $-40^{\circ}\text{C}$  before bubbles were measured. Heat-shock was achieved with 7 days cycle of temperature variation between  $-20$  and  $-8^{\circ}\text{C}$ . Each increase or decrease of  
20 temperature lasted 12 hours. Bubble size distribution was achieved by image analysis of the bubbles taken from 4 repetitions of the same sample after dispersion in a viscous glycerol containing medium. It is clear that the initial size distributions are very similar, which  
25 corroborates the previous microscopical observations on the bubbles (the sample produced with our invention required 2 times less energy input). After heat shock, both the control and BLG/AG control showed a shift of the size distribution to smaller air bubbles, which is in  
30 fact due to a loss of the largest air bubbles. Difference in air bubble content is the area between the initial and the heat-shock curves. Very interestingly, the curve

corresponding to our invention did not move at all. This reveals the very high heat-shock stability of the sample.

Figure 5 shows the confocal scanning laser microscopy pictures of the three ice-cream products after 7 days equilibration at  $-40^{\circ}\text{C}$ . The protein was coloured by the rhodamine 6G reagent that is known to specifically label protein through hydrophobic interactions. The confocal microscopy is carried out as follows : an aliquot of the product is put into a 2 mm deep covet and covered with 100  $\mu\text{l}$  of a  $10^{-6}$  M aqueous solution of Rhodamine 6G. After melting, the melting is covered with a glass cover slide and examined by confocal microscopy. On the figure, the views are following : (A) control recipe; (B) BLG/AG control recipe; (C) recipe according to our invention. From the observation of the pictures, the structuration of the air bubble interface with proteins using our invention is clear as compared to the two controls. In these two latter products, proteins are somewhat distributed randomly within the matrix.

Figure 6 represents the thin section micrographs of the three ice-cream products after Rhodamine 6G coloration according to the same procedure as for the preceding figure. On this figure, the views are following : (A) control recipe; (B) BLG/AG control recipe; (C) recipe according to hour invention. The presence of specific structures around the air bubbles has to be noted. To our opinion, these are coacervates made from the interfacial interaction between the primary formed BLG/AG complexes.

Figure 7 shows the confocal scanning laser micrograph of the interface of an air bubble made in a just made

mixture of  $\beta$ -lactoglobulin 2.5% and acacia gum 1.25% at pH 4.2 in water.  $\beta$ -lactoglobulin was covalently labeled with fluoresceine isothiocyanate. The coacervates present at the interface are comparable to those observed in the  
5 final ice cream product obtained according to our invention (Fig. 6C).

### Examples

10

The following examples illustrate some applications of the present invention.

#### Example 1

15

An concentrated emulsion is obtained preparing first an emulsion by mixing during 5 minutes using a mixer 0.74% whey protein isolate, 66% sunflower oil and water at pH 4.2 by addition of lactic acid. A second emulsion is then  
20 prepared by mixing during 5 minutes using a mixer 0.23% acacia gum powder, 66% sunflower oil and water at pH 4.2 by addition of lactic acid. The final concentrated emulsion is obtained by mixing the two previously prepared emulsions at a 1:1 weight ratio and mixing  
25 during 10 minutes using a mixer or a high shear pump.

#### Example 2

An ice cream mix is prepared from 9% palm-palm kernel  
30 oil, 5% non-fat milk solids, 5% whey protein isolate, 17% sucrose, 0.4% of a stabiliser blend containing hydrocolloids such as locust bean gum, guar, carrageenans, carboxymethylcellulose, water and

emulsifiers. The pH of this first mix is adjusted to pH 4.2 by addition of citric acid. A second mix is prepared from 9% palm kernel oil, 7.5% non-fat milk solids, 2.5% acacia gum powder, 14% sucrose, 0.4% of a stabiliser blend containing hydrocolloids such as locust bean gum, guar, carrageenans, carboxymethylcellulose, water and emulsifiers. The pH of this second mix is adjusted to pH 4.2 by addition of citric acid. Both mix are then homogenised at 100 bars using a homogeniser and then pasteurised. After maturation at 4°C, the two ice cream preparations were mix together at a 1:1 mixing ratio and aerated at -6°C to +4°C using the processing device described on Fig. 1 coupled with a shearing device (Hoyer freezer or Mondomix whipper) to obtain an overrun of 100%. The mix was then poured in molds and hardened at a temperature of -40°C.

### Example 3

A sour cream mousse is prepared from 50% milk cream (30% fat), 10% non-fat milk solids, 5% whey protein isolate, 8% sucrose, water and emulsifiers. The pH of this first recipe is adjusted to pH 4.3 by addition of citric acid. A second recipe is prepared from 50% milk cream (30% fat), 10% non-fat milk solids, 2.5% acacia gum powder, 8% sucrose, 0.4% of a stabiliser blend, water and emulsifiers. The pH of this second recipe is adjusted to pH 4.3 by addition of citric acid. Both preparations are then homogenised at 80 bars using a homogeniser and then pasteurised. The two preparations were mix together at a 1:1 mixing ratio and aerated at +4°C using the processing device described on Fig. 1 coupled with a Mondomix

whipper to obtain an overrun of 100%. The aerated sour cream mousse was then poured in molds and stored at +4°C.

#### Example 4

5

A wafer recipe is obtained by mixing using a mixer 50% of a wheat flour containing 70% of starch and 4% of a whey protein concentrate with water and adjusting the pH to 4.2 by addition of lactic acid. Another recipe is  
10 obtained by mixing using a Hobbard mixer 50% of a wheat flour containing 70% of starch and 1.6% of an acacia gum powder with water and adjusting the pH to 4.2 by addition of lactic acid. Both recipes are then mixed together at a 1:1 mixing ratio and aerated using the mixing device  
15 described on Fig. 1 coupled with Hobbard mixer equipped with a thermostated bowl. This aerated wafer base is then poured in-between baking plates at 150°C and cooked to obtain wafers.

#### 20 Example 5

A liquid dairy coffee creamer recipe is obtained by mixing and high-pressure homogenization (400 bars + 80 bars) 10% maltodextrin DE21, 15% hydrogenated palm oil,  
25 3% whey protein isolate with water at 50°C and adjusting the pH to 4.2 by addition of hydrochloric acid. Another recipe is obtained by mixing and high-pressure homogenization (400 bars + 80 bars) 10% maltodextrin DE21, 15% hydrogenated palm oil, 3% acacia gum powder  
30 with water at 50°C and adjusting the pH to 4.2 by addition of hydrochloric acid. Both recipes are then mixed together at a 1:1 mixing ratio using a high-shear mixer equipped with a thermostated bowl at 50°C.



**Claims**

- 5 1) A product taken from the group consisting of a foam,  
an emulsion, a foamed emulsion, a dispersed emulsion  
and a foamed dispersion, wherein the interface water-  
air, water-oil or water-solid comprises a complex  
formed instantaneously at said interface by the  
10 mixture of at least a protein (or peptide) and at  
least a polysaccharide oppositely charged or the  
mixture of two proteins oppositely charged, said  
product being in a pH range within which the  
electrostatic interaction between both compounds  
15 oppositely charged occurs and wherein the total amount  
of protein and polysaccharide is comprised between  
0.01 and 5 % in weight.
- 20 2) A product according to claim 1, wherein the protein is  
taken from the group consisting of a protein from  
milk, soy, egg, meat, fish and plant.
- 25 3) A product according to claim 2, wherein the protein is  
taken from the group consisting of  $\beta$ -lactoglobulin ,  
whey protein, soy protein, egg white protein and wheat  
protein.
- 30 4) A product according to any of claims 1 to 3, wherein  
the polysaccharide is taken from the group consisting  
of gums, hydrocolloids and stabilizers.
- 5) A product according to claim 4, wherein the  
polysaccharide is acacia gum , arabinogalactans, Ax

rye, Ax wheat, carboxy-methyl-cellulose, chitosan, xanthan gum,  $\beta$ -glucan.



- 5 6) A product according to any of claims 1 to 5, wherein the ratio protein to polysaccharide or protein to protein is comprised between 1:20 and 20:1.
- 10 7) The use of the product according to any of claims 1 to 6, in the ice-cream, low-fat mayonnaise, culinary, chocolate, dessert, wafers, sponge cakes, dairy and petfood area, wherein said product is used in an amount comprised between 10 and 100 % of the final product in weight.
- 15 8) The use of the product according to any of claims 1 to 6 in the cosmetic and perfume area, wherein said product is used in an amount comprised between 10 and 100 % of the final product in weight.
- 20 9) A process for the preparation of a foam product, wherein a solution or a bulk mass of the at least one protein and a solution or a bulk mass of the at least one polysaccharide or a solution or a bulk mass of one protein and another solution or a bulk mass of one protein is injected with the air in a bulk mass or
- 25 directly in the air.
- 30 10) A process for the preparation of a foam product, wherein a solution of at least one protein and a solution of at least one polysaccharide or a solution of one protein and another solution of one protein are mixed together in the presence of air.

- 11) A process for the preparation of an emulsion, wherein a first part of an emulsion is stabilised with the at least one protein, a second part of an emulsion is stabilised with the at least one polysaccharide or a second protein, and both emulsions are mixed together.
- 12) A process for the preparation of a foamed emulsion, wherein a bulk product is prepared, a first part of an emulsion is stabilised with at least one protein, a second part of an emulsion is stabilised with at least one polysaccharide or a second protein, both emulsions are mixed together and diluted in the bulk product, then a new solution of the protein with a new solution of the polysaccharide are injected with air in the bulk product to form the foamed bulk product.
- 13) A process for the preparation of a dispersed emulsion, wherein a first part of an emulsion is stabilised with at least one protein, a second part of an emulsion is stabilised with the at least one polysaccharide or a second protein, and both emulsions are mixed together, the obtained final emulsion being then mixed with a base comprising particles.
- 14) A process for the preparation of a foamed dispersion, wherein a solution of the at least one protein and a solution of the at least one polysaccharide or a solution of one protein and another solution of one protein is injected with the air in a bulk of dispersed particles.
- 15) A device for carrying out the process according to any of claims 9 to 14, which comprises on a frame :

- a first pipe for the arrival of an emulsion or dispersion with a protein
  - a second pipe for the arrival of the gas
  - a third pipe for the arrival of an emulsion or
- 5 dispersion with a polysaccharide,

these three pipes arriving on a main channel, being disposed perpendicular and staggered along said main channel , the first pipe forming the central pipe on the main channel, the second pipe forming the intermediate

10 pipe on the main channel and the third pipe forming the external pipe on the main channel and wherein the outlet on the main channel of the central and the intermediate pipe are staggered.








Fig. 3




Fig. 4




Fig. 5






Fig. 6




Fig. 7

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 03/10740

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A23P1/16 A23L1/05 A23J3/00 A23G3/00 A23G1/00  
A21D13/00 A23L1/053 A23L1/054 A23L1/056 A23L1/0534  
A61K7/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A23P A23L A23J A23G A21D A61K A23D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, FSTA

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	POOLE S: "The foam-enhancing properties of basic biopolymers." INTERNATIONAL JOURNAL OF FOOD SCIENCE & TECHNOLOGY 1989, vol. 24, no. 2, 1989, pages 121-137, XP008014428 UK cited in the application the whole document	1-14
X	GB 2 134 117 A (BRITISH FOOD MANUFACTURING IND) 8 August 1984 (1984-08-08) cited in the application the whole document	1-14

-/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

22 January 2004

Date of mailing of the international search report

09/02/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Vuillamy, V



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/10740

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB 2 179 043 A (BRITISH FOOD MANUFACTURING IND) 25 February 1987 (1987-02-25) cited in the application the whole document -----	1-14
X	AHMED M ET AL: "Foaming of aqueous solutions of protein + propylene glycol alginate." FOOD HYDROCOLLOIDS 1991, vol. 4, no. 5, 1991, pages 395-402, XP008014429 UK cited in the application the whole document -----	1-14
X	US 5 342 643 A (WOLF PETER A ET AL) 30 August 1994 (1994-08-30) the whole document -----	1-8,11
X	PATENT ABSTRACTS OF JAPAN vol. 0091, no. 30 (C-284), 5 June 1985 (1985-06-05) & JP 60 016573 A (RAION KK), 28 January 1985 (1985-01-28) abstract -----	1-8,11
X	GIRARD M ET AL: "Emulsifying properties of whey protein-carboxymethylcellulose complexes." JOURNAL OF FOOD SCIENCE, vol. 67, no. 1, 2002, pages 113-119, XP002267658 the whole document -----	1-8,11
X	EINHORN-STOLL U ET AL: "MODIFIED PECTINS IN WHEY PROTEIN EMULSIONS" DIE NAHRUNG, vol. 40, no. 2, 1996, pages 60-67, XP000604246 DE ISSN: 0027-769X the whole document -----	1-8,11
X	GARTI N ET AL: "Portulaca oleracea gum and casein interactions and emulsion stability." FOOD HYDROCOLLOIDS, vol. 13, no. 2, 1999, pages 127-138, XP002267659 the whole document -----	1-8,11

-/--

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/10740

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DICKINSON E ET AL: "Influence of kappa-carrageenan on the properties of a protein-stabilized emulsion." FOOD HYDROCOLLOIDS, vol. 12, no. 4, 1998, pages 417-423, XP002267660 the whole document	1-8, 11
X	DICKINSON E: "Stability and rheological implications of electrostatic milk protein-polysaccharide interactions." TRENDS IN FOOD SCIENCE & TECHNOLOGY, vol. 9, no. 10, 1998, page 347354, XP002267661 the whole document	1-8, 11
X	GB 1 440 181 A (UNILEVER LTD) 23 June 1976 (1976-06-23) the whole document	1-8, 11
X	WO 00/23038 A (REVLON CONSUMER PROD CORP) 27 April 2000 (2000-04-27) the whole document	1-8, 11, 13
X	& US 6 197 319 B1 (RUSS JULIO GANS ET AL) 6 March 2001 (2001-03-06) cited in the application	1-8, 11, 13
X	GB 751 600 A (NCR CO) 27 June 1956 (1956-06-27) the whole document	1-8, 11
X	CH 675 370 A (CIBA GEIGY AG) 28 September 1990 (1990-09-28) figure 5	15

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/EP 03/10740

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-8 partly, 9, 10, 12, 14, 15

Product, process and device for stabilizing a product with a protein-polysaccharide or protein-protein complex, where the product is a foam, a foamed emulsion or a foamed dispersion.

---

2. claims: 1-8 partly, 11, 13

Product and process for stabilizing a product with a protein-polysaccharide or protein-protein complex, where the product is an emulsion or a dispersed emulsion.

---

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 03/10740

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 2134117	A	08-08-1984	NONE	
GB 2179043	A	25-02-1987	NONE	
US 5342643	A	30-08-1994	NONE	
JP 60016573	A	28-01-1985	NONE	
GB 1440181	A	23-06-1976	AT 332203 B AT 431673 A AU 5575873 A BE 799772 A1 BE 799773 A1 BE 506697 A CA 1018821 A1 CH 582483 A5 DE 2325131 A1 DE 2325133 A1 ES 414862 A1 FI 55287 B FR 2185018 A1 FR 2185020 A1 FR 1048667 A IE 37634 B1 IT 500010 A IT 991590 B JP 49066847 A NL 7306716 A ,B NL 7306863 A NL 7309962 A NL 7314932 A ,B SE 390868 B ZA 7303238 A	10-09-1976 15-12-1975 21-11-1974 19-11-1973 19-11-1973  11-10-1977 15-12-1976 17-01-1974 13-12-1973 16-06-1976 30-03-1979 28-12-1973 28-12-1973 23-12-1953 14-09-1977  30-08-1975 28-06-1974 20-11-1973 20-11-1973 21-01-1974 02-05-1974 31-01-1977 29-01-1975
WO 0023038	A	27-04-2000	AU 1218500 A WO 0023038 A1 US 6197319 B1	08-05-2000 27-04-2000 06-03-2001
US 6197319	B1	06-03-2001	AU 1218500 A WO 0023038 A1	08-05-2000 27-04-2000
GB 751600	A	27-06-1956	US 2800458 A BE 530008 A BE 530009 A BE 530010 A CH 327303 A CH 330500 A CH 334334 A DE 1010822 B DE 1017905 B FR 1114850 A FR 1114851 A FR 1165805 A GB 759800 A GB 750972 A GB 873943 A NL 95043 C NL 95044 C	23-07-1957    31-01-1958 15-06-1958 30-11-1958 19-06-1957 17-10-1957 17-04-1956 17-04-1956 29-10-1958 24-10-1956 20-06-1956   

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Publication No

PCT/EP 03/10740

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 751600	A	NL 95045 C US RE24899 E	
CH 675370	A	CH 675370 A5	28-09-1990